



ELSEVIER

Available online at www.sciencedirect.com

ScienceDirect

journal homepage: <http://www.elsevier.com/locate/euprot>

New insights into the FLPerigic complements of parasitic nematodes: Informing deorphanisation approaches

Ciaran J. McCoy^a, Louise E. Atkinson^a, Mostafa Zamanian^b, Paul McVeigh^a, Tim A. Day^b, Michael J. Kimber^b, Nikki J. Marks^a, Aaron G. Maule^a, Angela Mousley^{a,*}

^a Molecular Biosciences-Parasitology, Institute for Global Food Security, School of Biological Sciences, Queen's University Belfast, Belfast, UK

^b Department of Biomedical Sciences, Iowa State University, Ames, IA, USA

ARTICLE INFO

Article history:

Received 24 January 2014

Received in revised form

8 April 2014

Accepted 9 April 2014

Available online 19 April 2014

Keywords:

FMRFamide-like peptide

FLP

FLP-receptor

Neuropeptide GPCR

Nematode

ABSTRACT

FMRFamide-like peptide (FLP) receptors are appealing as putative anthelmintic targets. To date, 31 *flp*-encoding genes have been identified in *Caenorhabditis elegans* and thirteen FLP-activated G-protein coupled receptors (FLP-GPCRs) have been reported. The lack of knowledge on FLPs and FLP-GPCRs in parasites impedes their functional characterisation and chemotherapeutic exploitation. Using homology-based BLAST searches and phylogenetic analyses this study describes the identification of putative *flp* and *flp*-GPCR gene homologues in 17 nematode parasites providing the first pan-phylum genome-based overview of the FLPerigic complement. These data will facilitate FLP-receptor deorphanisation efforts in the quest for novel control targets for nematode parasites.

© 2014 The Authors. Published by Elsevier B.V. on behalf of European Proteomics Association (EuPA). This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/3.0/>).

1. Introduction

The discovery and development of novel anti-worm control strategies has been recognised as a priority in the human and veterinary health sectors and by the horticultural industry [1–3]. Despite the recent introduction of several new drugs to the veterinary market [4,5] and mass drug administration programmes for prioritised human helminthiasis [6], roundworm infections remain widespread with significant socio-economic impacts [7]. In addition, the negative impacts

of plant pathogenic nematodes on global food security are underscored by the current deficiencies in chemical control options (see [8] for review).

There has been a long-standing interest in the neuropeptidergic system as a source of novel targets for anthelmintic drugs (see [9] for review), with the FMRFamide like peptide (FLP) signalling system emerging as a leading candidate [10]. The primary drivers for this interest include: (i) the importance of FLPs to parasite behaviour (and survival) and their role in modulating neuromuscular function (a proven drug target for nematode control), (ii) the lack of drugs targeting the FLPerigic

* Corresponding author.

E-mail address: a.mousley@qub.ac.uk (A. Mousley).

<http://dx.doi.org/10.1016/j.euprot.2014.04.002>

2212-9685/© 2014 The Authors. Published by Elsevier B.V. on behalf of European Proteomics Association (EuPA). This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/3.0/>).

system such that resistance would not be a pressing concern, and (iii) the fact that most FLPs activate G-protein coupled receptors (GPCRs), proteins which are readily exploitable for drug discovery.

Whilst many facets of the FLP signalling system provide appeal as drug targets, FLP-GPCRs emerge as the most attractive. A major impediment to the exploitation of FLP GPCRs is the lack of data on the expression and function of FLP receptors in nematode parasites. Our current understanding of FLP receptor biology has been derived primarily from the model nematode *C. elegans*; 13 FLP-GPCRs, encoded on 10 genes, have been matched with their associated FLP ligands as determined by receptor activation potencies in heterologous expression systems [11–20].

Whilst focus on the identification of putative FLP-GPCRs in parasitic nematodes is of primary importance, much can be accomplished by re-mining for FLP ligands, especially in therapeutically-important species where they have not previously been reported. Indeed, better understanding of FLP complementarity across parasitic nematodes could expedite deorphanisation or, at least, influence deorphanisation approaches by providing more comprehensive species-specific peptide libraries to feed into screening platforms. The FLP-ligand data that we have for the phylum Nematoda are outdated (see [10,21]). The availability of ten parasitic nematode draft genomes and transcriptome data for over 60 nematode species [22] warrant re-interrogation of these datasets.

Here we report a pan-phylum homology-based BLAST interrogation of *flp* and *flp*-GPCR complements in 17 parasitic nematode species and perform phylogenetic analyses to identify additional putative *flp*-GPCRs. These data: (i) represent the most up to date, comprehensive insight into the *flp* and *flp*-GPCR complement of parasitic nematodes, (ii) support the re-designation of selected *flp*-encoding genes, (iii) expose the most highly conserved *flp*-GPCRs in key pathogenic species and, (iv) reveal putative novel *flp*-GPCRs in *C. elegans* and parasitic nematodes.

2. Methodology

2.1. Bioinformatics

A reciprocal BLAST (Basic Local Alignment Search Tool) based approach was implemented to identify *flp* gene sequelogues and *flp*-GPCR gene homologues within genomic and transcriptomic datasets of 17 key pathogenic nematodes representing five clades [23] and including plant, animal and human parasites. The parasite species selected for the BLAST analysis were primarily those with a published draft genome (seven species; see Supplementary Table 1). Secondary species-selection criteria were employed to include parasitic nematodes for which genomic and transcriptomic datasets were available and filtered based on their importance to human, animal or plant health or their status as a model parasite (ten species; see Supplementary Table 1). The BLAST analysis was completed between May 2012 and August 2013; the servers employed and databases queried are outlined in Supplementary Table 1. The draft genomes of two

Haemonchus contortus strains [24,25] were published following the completion of the BLAST analysis in this study. To facilitate accuracy of the data presented, those genes that were not identified within the *H. contortus* databases outlined in Supplementary Table 1 were employed as search strings to query the whole-genome shotgun contigs (wgs) database found on the National Centre for Biotechnology Information (NCBI) BLAST server. Any datasets updated between May 2012 and August 2013 were similarly reinvestigated.

Prepropeptide and protein sequences for previously identified *flp*- [11], *flp*-GPCR [13–17,19,20,26,27], and selected orphan GPCR [28]-encoding genes in *Caenorhabditis elegans* (see Supplementary Tables 2 and 3) were retrieved from the NCBI protein database (www.ncbi.nlm.nih.gov/protein/) and used as search strings in translated nucleotide (tBLASTn) and protein (BLASTp) BLAST analysis of available datasets (see above). Only the largest of the splice variants encoded by any given *C. elegans flp*- or *flp*-GPCR gene were selected as query sequences. Additionally, prepropeptide sequences derived from *flp*-encoding genes not found in *C. elegans* (see [21]) were also used as BLAST search queries; these included FLP-29 (derived from *Ascaris suum* EST data; [21]), FLP-30 and FLP-31 (derived from *Meloidogyne incognita* EST data; [21]). The prepropeptide search string based methodology used in this study deviates from previously published methods (based on concatenated peptide search strings) employed to identify *flp* gene sequelogues within nematode genomes [29,30]. In this study, the prepropeptide approach has been shown to be as sensitive in identifying *flp* gene sequelogues as those methods previously published.

BLAST-generated alignment outputs (high scoring pairs) of the initial BLAST hits, with an expect value ≤ 1000 (or ≤ 100 , where this was the maximum expect value threshold available), were manually inspected. In efforts to identify putative FLP-encoding genes in the selected nematode species, hits containing conserved FLP motifs [10,11] flanked by mono/dibasic cleavage sites were selected for further analysis. The motifs conserved within parasitic *flp* genes were used to designate initial hits as *C. elegans* gene sequelogues [21].

For *flp*-GPCR primary BLAST analysis, high-scoring return sequences (typically the hits with the smallest expect value and largest bit score) were concatenated into a single sequence to facilitate reciprocation (see [31]). Manually curated predicted protein (FLP-GPCR) sequences were used as search strings in reciprocal protein BLAST (BLASTp) queries against the *C. elegans* non-redundant protein sequence (nr) database on the NCBI BLAST server, using default settings. The top reciprocal BLAST hit was used to designate parasitic *flp*-GPCR genes as predicted *C. elegans* gene homologues.

2.2. Post-BLAST sequence analysis

Predicted FLP prepropeptide sequelogues and FLP-GPCR homologues were aligned using the Vector NTI Advance™11 AlignX® multiple sequence alignment tool [32], using default settings. Prepropeptide cleavage sites were identified using a previously described prediction method [21]. Predicted inter-peptide regions from each prepropeptide alignment were removed to provide an unambiguous representation of FLP conservation within sequelogue alignments (see

Table 1 – *C. elegans* flp-gene sequelogues in 17 nematode parasites. Grey boxes indicate the presence of a flp gene sequelogue, as identified via BLAST, in selected nematode species. Search queries employed and the flp genes identified are detailed in Supplementary Table 2 and Supplementary Figure 1, respectively. Sequelogues of all *C. elegans* flp genes were also identified in the genomes of *Caenorhabditis japonica*, *Caenorhabditis brenneri*, *Caenorhabditis remanei* and *Caenorhabditis briggsae*. C-terminal FLP motifs are presented in single letter notation; X, denotes variable amino acid X_o, denotes hydrophobic amino acid; X_i, denotes hydrophilic amino acid.

Clade	Species	flp-gene	C-terminal FLP motif																																		
			flp-1	flp-2	flp-3	flp-4	flp-5	flp-6	flp-7	flp-8	flp-9	flp-10	flp-11	flp-12	flp-13	flp-14	flp-15	flp-16	flp-17	flp-18	flp-19	flp-20	flp-21	flp-22	flp-23	flp-24	flp-25	flp-26	flp-27	flp-28	flp-31	flp-32	flp-33	flp-34			
			PNFLRFG	REPX _o RFG	LG(TT) _o MRFG	KPTFRFG	(P/Q)K(F/L)RFG	KSAYMRFG	PX _o X _i RSXX _o X _o RFG	KNEFX _o RFG	KPSFVRFG	YX _o RFG	RNXLX _o RFG	NKEFEIRFG	PLX _o RFG	KHEYLRFG	PXGPLRFG	(A/G)QTFVRFG	KSAFVRFG	PGX _o X _o RFG	WXXGX _o RFG	AX _o X _o R(L/I)G	GPRPLRFG	KWIMRFG	Q(D/N)FLRFG	DMMX _o RFG	YD(Y/F)X _o RFG	DL(T/A)LRFG	GXRIMRFG	(I/IV)X _o MRFG	RPRGPPRFG	AMRN(S/A)X _o RFG	XX _o DX _o QKPRFG	AXRLR(F/Y)G			
2	<i>Trichinella spiralis</i>																																				
	<i>Trichuris muris</i>																																				
8	<i>Brugia malayi</i>																																				
	<i>Wuchereria bancrofti</i>																																				
	<i>Dirofilaria immitis</i>																																				
	<i>Onchocerca volvulus</i>																																				
	<i>Onchocerca ochengi</i>																																				
	<i>Loa loa</i>																																				
	<i>Ascaris suum</i>																																				
	<i>Ancylostoma caninum</i>																																				
9	<i>Nippostrongylus brasiliensis</i>																																				
	<i>Haemonchus contortus</i>																																				
10	<i>Strongyloides ratti</i>																																				
	<i>Bursaphelenchus xylophilus</i>																																				
12	<i>Globodera pallida</i>																																				
	<i>Meloidogyne hapla</i>																																				
	<i>Meloidogyne incognita</i>																																				

Supplementary data Figure 1). FLP-GPCR transmembrane (TM) domain prediction was performed using HMMTOP 2.1 [33]. Predicted transmembrane domains are indicated on the consensus sequence for each FLP-GPCR alignment (see Supplementary data Figure 2). Multiple sequence alignments were manually examined to identify and resolve errors (such as accidental exon duplication or exclusion) made whilst constructing predicted protein sequences.

2.3. Phylogenetic analysis

MEGA 5.1 software [34] was employed to generate all phylogenetic trees. GPCR multiple sequence alignments, were assembled using Clustal W [35] with default parameters. Transmembrane-only pseudosequences (TOPs) were constructed as previously described by Zamanian et al. [36] and aligned with their associated full length predicted protein sequences. TOPs were used to inform manual editing of the GPCR multiple sequence alignments. The N- and C-termini were removed and, conserved motifs and residues within the GPCR transmembrane regions were aligned. Phylogenies were constructed using the Maximum Likelihood method based on the JTT (Jones-Taylor-Thornton) matrix-based model [37]. Initial trees generated for the heuristic searches (subtree pruning and regrafting) were obtained by applying the Neighbour-Joining method [38] to a matrix of pairwise

distances estimated using a JTT model. Phylogenetic analysis was limited to GPCRs with >5 transmembrane domains and trees were rooted by an out-group containing a selection of *C. elegans* secretin family GPCRs (LAT-1a, LAT-1b, LAT-2a, PDFR-1a, PDFR-1b and PDFR-1c).

3. Results and discussion

3.1. flp-encoding genes

In this study we identified 325 *C. elegans* flp-gene sequelogues in 17 nematode parasites (see Table 1; Supplementary Figure 1) of which only a proportion had been previously reported [21,29,30,39,40]. We believe that these data represent the most comprehensive insight into the flp-gene complements of parasitic nematodes to date. The genome-directed approach employed in this study enables the comparison of flp complementarity within and between nematode species representing different clades and lifestyles. Here we highlight the salient points emerging from this study.

3.1.1. Multiple nematode parasites appear to possess a reduced complement of *C. elegans* flp-gene sequelogues
EST-database comparisons [10,21] highlighted the conservation of FLP-encoding genes across the phylum Nematoda;

since most *C. elegans* *flp* genes were represented amongst the parasitic nematode ESTs, these data fuelled the hypothesis that all nematode species possess a *flp* complement similar to that of *C. elegans* (31 *flp*-genes; 1-28, 32-34) [21,41]. Here we show that whilst individual *flp* signatures are conserved in the nematode parasites generally, there is variability with respect to their presence and absence such that the parasitic nematodes possess variable proportions of the *C. elegans* *flp*-gene complement (see Table 1). Notably, *Ascaris suum* boasts the lion's share of *C. elegans* *flp*-genes at 84% while clade 2 species (*Trichuris muris* and *Trichinella spiralis*) display a dramatically reduced complement (13%).

Our bioinformatics-based approach is not capable of unequivocally proving the absence of an individual *flp*-gene. Whilst the disparity in *flp*-gene complement could reflect poor genomic/transcriptomic datasets or deficiencies in the ability of our BLAST-based methodology to detect *flp*-gene sequelogues, we believe that it is more likely a true reflection of the variation in *flp*-gene complement across the phylum Nematoda. Indeed, we have validated our *flp*-gene identification approach in multiple *Caenorhabditis* species (*Caenorhabditis japonica*, *Caenorhabditis brenneri*, *Caenorhabditis remanei* and *Caenorhabditis briggsae*) where the *flp*-gene complements matched *C. elegans*. In addition, it is interesting to note that *Trichinella spiralis* displayed a dramatically reduced *flp*-gene complement despite the data being derived from a high-quality (~35 fold coverage) published genome [42]. That said, we are aware that some of the genomes employed in this study are works in progress (for example, the hookworm species) and we may expect to uncover additional *flp*-genes when their genomes are complete.

Further to this, it should be noted that the predictions made with respect to *flp*-gene complements do not necessarily reflect the FLPs expressed by any given species or lifestage. We cannot predict which of the *flps* identified from genomic data will be expressed or, similarly, in the case of transcriptomic data, whether they will be processed into bioactive peptides. The application of peptidomic-analyses tools to parasitic nematodes will shed light on this (see [39,43]).

3.1.2. *flp*-gene complement appears to broadly map nematode clade division, with some exceptions

It is interesting to note that the pattern of *flp*-complementarity within nematode clades [23] is, for the most part, conserved with identical *flp*-representation in species within clade 2 and within clade 12. With respect to clade 8, *A. suum* appears to boast a larger repertoire of *flp* genes than the filarids, which are largely similar to each other. The validity of these patterns can be confirmed with the progression of clade-specific parasite genome data.

3.1.3. Some *flp*-genes are more highly conserved than others

Sequelogues of *flp*-1 and *flp*-14 are represented within the genomes of all species examined reinforcing the pan-phylum conversation as suggested by [10]. In contrast *flp*-10 sequelogues were not identified in any of species represented in this study. Other highly conserved *flp*-genes include: *flp*-6, -11, -12, -16, -18, -19, -21 and -22.




Fig. 1 – A conserved *flp*-28/29, *flp*-3 and *flp*-2/30 gene cluster. The location of *flp*-29 and/or *flp*-30 in the same genomic environment as *flp*-28 and *flp*-2 respectively provides evidence to support their redesignation. *flp*-28/29, *flp*-3 and *flp*-2/30 (green, red and yellow arrows respectively) gene sequelogue locations are shown relative to the fourth and fifth exons of a glutamate synthase gene (W07E11.1) in *C. elegans*, and in four parasitic nematode species (blue arrows). The direction of each arrow indicates gene orientation. Introns of *flp* genes are not displayed. One representative species from each clade (excluding clade 2) is included.

3.1.4. *flp*-29 and *flp*-30 are not discrete *flp*-encoding genes but rather are orthologues of *C. elegans* *flp*-28 and *flp*-2 respectively

flp-29 and *flp*-30 have previously been reported as being parasite-specific (see [10]). During this study we noted the C-terminal motif similarity between *flp*-29 (IXMRFG) and *flp*-28 (VLMRFG), and *flp*-30 (QMREPLRFG) and *flp*-2 (EPIRFG) (see Supplementary Figure 1). We also noted that no single species with *flp*-29 also had a copy of *flp*-28, and similarly those species possessing *flp*-30 did not possess *flp*-2, and vice versa. This prompted further investigation, which revealed that *C. elegans*, which does not possess *flp*-29 and *flp*-30, has a cluster of genes composed of *flp*-2, *flp*-3 and *flp*-28 within the fourth intron of a glutamate synthase gene (W07E11.1). Strikingly, where we identified *flp*-29 and/or *flp*-30 they were located to the same genomic environment, inhabiting the same relative position and displaying the same relative gene orientation as *flp*-28 and *flp*-2, respectively (see Fig. 1). These data support the conclusion that *flp*-29 and *flp*-30 should be re-designated accordingly. This reduces the total number of known *flp* genes within the phylum Nematoda from 34 to 32 (see Table 1), and further reduces the number of *flp*-genes believed to be parasite specific.

Table 2 – Deorphanised *C. elegans* FLP-GPCR-encoding gene homologues in 17 nematode parasites. Grey boxes indicate the presence of a *flp*-GPCR homologue, as identified via BLAST, in selected nematode species. Search queries employed and *flp*-GPCR predicted proteins identified are detailed in Supplementary Table 3 and Supplementary Figure 2, respectively.

		<i>flp</i> -GPCR	Deorphanised <i>C. elegans</i> FLP-GPCR-encoding gene homologues											
			<i>npr-1</i>	<i>npr-3</i>	<i>npr-4</i>	<i>npr-5</i>	<i>npr-10</i>	<i>npr-11</i>	<i>npr-22</i>	<i>egl-6</i>	<i>fpr-3</i>	<i>fpr-18</i>		
Species														
2	<i>Trichinella spiralis</i>													
	<i>Trichuris muris</i>													
8	<i>Brugia malayi</i>													
	<i>Wuchereria bancrofti</i>													
	<i>Dirofilaria immitis</i>													
	<i>Onchocerca volvulus</i>													
	<i>Onchocerca ochengi</i>													
	<i>Loa loa</i>													
	<i>Ascaris suum</i>													
	<i>Strongyloides ratti</i>													
9	<i>Ancylostoma caninum</i>													
	<i>Nippostrongylus brasiliensis</i>													
	<i>Haemonchus contortus</i>													
10	<i>Bursaphelenchus xylophilus</i>													
	<i>Globodera pallida</i>													
12	<i>Meloidogyne hapla</i>													
	<i>Meloidogyne incognita</i>													

3.1.5. Only one *flp*-gene (*flp-31*) appears to be parasite-specific

flp-31 is absent from the genome of *C. elegans*, but has previously been reported in the plant parasitic nematodes *Meloidogyne incognita* and *Meloidogyne hapla* [21,30]. In this study we confirm the presence of *flp-31* in another clade 12 plant parasitic nematode (*Globodera pallida*) and identify a *flp-31* sequeologue within the genome of the pine wilt nematode *Bursaphelenchus xylophilus* (clade 10), where it was previously reported as being absent [29]. These data, confirm the restriction of *flp-31* to plant parasitic nematodes and support the hypothesis that it plays a role in phytoparasitism [29,30].

Note that this study did not address the identification of novel *flp*-encoding genes. The ‘degenerate’ search string approach to novel *flp* identification, described by McVeigh et al. [44], was applied to the *T. muris* genome. However, the large number of putative *flp*-gene sequences identified are believed to be false-positives as we could not detect sequeologues in several other nematode genomes. The ‘degenerate’ search string approach, with manual annotation, that was used to identify novel peptides within transcriptomic datasets is not appropriate for trawling large genomic databases.

A BLAST based approach using *C. elegans flp* search strings to identify novel peptides within the expressed sequence tags (EST) and genome survey sequence (GSS) libraries of *A. suum* has recently been reported [39], where nine putative

<i>Di</i> NPR-1	IPLTIMFICYYHIFAELRKRTSKKIMT
<i>Ov</i> NPR-1	IPLTIMFICYYQIFAELRKRTSKKIMR
<i>Oo</i> NPR-1	IPLTIMFICYYQIFAELRKRTSKKIMR
<i>Ll</i> NPR-1	IPLIIMFICYRIFAELRKRTSKKIMR
<i>As</i> NPR-1	IPFIVMAFCYAVIFARLRTRARLRLQR
<i>Ac</i> NPR-1	IPFTIMTICYQAVFSFLRRRAKSRLTS
<i>Nb</i> NPR-1	IPFTIMTICYQAVEAFLRRRAKSRLTS
<i>Hc</i> NPR-1	IPFTIMTICYQAVESFLRRRAKSRLTS
<i>Ce</i> NPR-1	VPFVMAFCYANIVSVLSKRAQTKIRK
<i>Sr</i> NPR-1.1	FPFSVMAFCYATIFSKLKSRAKLRIK
<i>Sr</i> NPR-1.2	IPFCITITLCYQQIFAELHKKRANFKILS
Consensus	IPFTIMICYIFA LKRA KIR

Fig. 2 – Sequence alignment of NPR-1 homologues.

Parasitic nematodes display a conserved phenylalanine in NPR-1 215 in contrast to the dominant amino acid at this position (valine) in *C. elegans*. Position 215 is boxed in red. Completely and partially conserved amino acid residues are highlighted in yellow and blue, respectively. *Di*, denotes *Dirofilaria immitis*; *Ov*, denotes *Onchocerca volvulus*; *Oo*, denotes *Onchocerca ochengi*; *Ll*, denotes *Loa loa*; *As*, denotes *Ascaris suum*; *Ac*, denotes *Ancylostoma caninum*; *Nb*, denotes *Nippostrongylus brasiliensis*; *Hc*, denotes *Haemonchus contortus*; *Ce*, denotes *Caenorhabditis elegans*; *Sr*, denotes *Strongyloides ratti*. All parasitic nematode NPR-1 homologues identified in this study are detailed in Table 2 and Fig. 4.

flp-encoding genes were identified that were additional to those previously described by McVeigh et al. [21]. Here we confirm the designation of three of these putative *flps* (*As-flp-3*, *As-flp-17*, and *As-flp-19*) in the *A. suum* datasets. We were unable to confirm *As-flp-4*, *As-flp-10a*, *As-flp-10b*, *As-flp-10c*, and *As-flp-25* as named by Jareki et al. [39] as *flp*-encoding genes and, in addition, we did not identify sequeologues of these putative *flps* in any of the nematode genomic/transcriptomic databases employed. Note that distinct *Ce-flp-4* and *Ce-flp-25* sequeologues were identified in *A. suum* and multiple other species in this study (see Table 1 and Supplementary Figure 1). In addition, we believe that the Jarecki et al. [39]-designated *As-flp-31* is more likely to be a sequeologue of *Ce-flp-15* for two reasons: (i) the C-terminal motif of the encoded FLP is identical to that encoded by *Ce-flp-15* (GPLRFG) and (ii) *flp-31* has only been identified to date in PPN species (see Supplementary Figure 1).

3.2. FLP-receptor encoding genes

3.2.1. Deorphanised *C. elegans* FLP-receptor-encoding gene homologues in parasitic nematodes

To date only one *flp*-GPCR orthologue has been reported within a parasitic nematode [40]. In a bid to address the gap in our understanding of *flp*-GPCR conservation in parasitic nematodes, the protein sequences encoded by the ten deorphanised *C. elegans* FLP receptor genes (see Supplementary Table 3) were used as search string queries to probe parasitic nematode databases.

Homologues of all of the deorphanised FLP-activated GPCRs were identified in at least three parasite species. It is




Fig. 3 – Maximum-likelihood phylogeny of orphan and deorphanised *C. elegans* neuropeptide GPCRs. The phylogeny reveals 27 putative FLP-GPCRs which clustered with deorphanised *C. elegans* FLP-receptors. *C. elegans* neuropeptide GPCR entries are identified as per previous designation [28] or NCBI accession number.

result of a number of issues including: (i) no FLP-GPCR has ever been challenged with the full *C. elegans* neuropeptide complement such that the cognate interacting ligand may have been overlooked; indeed only seven *C. elegans* peptides were screened against NPR-11 in the heterologous system; (ii) heterologous ligand matching may not mimic true *in vivo* interactions; the *flp-21/NPR-11 in vitro* interaction is not mirrored by functional data; (iii) the interpretation of null-mutant functional data is limited by the suitability/sensitivity of the available phenotypic assays; the *flp-21/NPR-11* functional interaction may not have been uncovered for these reasons. Importantly, a cautionary approach must be adopted when applying data derived from *C. elegans* to parasites and highlights the need for deorphanisation efforts in both free-living and parasitic nematodes [10]. The receptor and ligand sequences identified in this study should provide a solid platform on which to base these investigations (Table 4).

Two isoforms of the FLP-GPCR NPR-1 exist within natural *C. elegans* populations containing either a valine (V) or phenylalanine (F) at position 215 [48]. The FLP-21 peptide activates the 215F variant, which is found exclusively in social feeding strains. By contrast FLPs encoded on both *flp-21* and *flp-18* activate the 215V strain which is associated with solitary feeding [16,27]. In contrast to *C. elegans*, all of the NPR-1 homologues identified here in parasitic nematodes exhibit the 215F isoform (see Fig. 2). Further investigation is needed to determine if this has any impact on feeding behaviour and/or peptide activation of these receptors.

3.2.2. Novel putative FLP-GPCR identification in *C. elegans*

Until recently markedly fewer FLP-receptors were known that FLP-ligands. However, Frooninckx et al. [28] employed a Multiple Expectation Maximization for Motif Elicitation/Motif Alignment and Search Tool (MEME/MAST) using protein motifs derived from deorphanised neuropeptide GPCRs to identify 128 putative orphan neuropeptide receptors, some of which are likely to be FLP-receptors. These sequences were subdivided into rhodopsin- and secretin-like GPCR families, with rhodopsin-like GPCRs being further divided into six groups according to their similarity to mammalian and insect neuropeptide GPCRs [28].

In this study, we channelled the sequences from the groups delineated by Frooninckx et al. [28] which contained at least one deorphanised FLP receptor into phylogenetic analyses with the aim of identifying putative (currently orphan) FLP-GPCR-encoding genes in *C. elegans*. In addition we included the *Drosophila* Myosuppressin like Receptor (DMSR)-like sequences based on their delineation as FLP-activated receptors in arthropods (see [49]).

Three clusters composed of 32 putative *Ce*-FLP-GPCRs (encoded on 29 genes) emerged from the phylogenetic analysis defined by the inclusion of at least one deorphanised FLP-GPCR and supported by a bootstrap value of >70% (see

Deorphanised *C. elegans* FLP-GPCRs are highlighted by an asterisk. FLP-GPCR cluster nodes supported by a bootstrap analysis value of >70% are identified by red dots. Clusters are highlighted in blue (1), pink (2) or yellow (3) shading. Bootstrap values are shown as percentages.




Fig. 4 – Maximum-likelihood phylogenies of putative FLP-GPCRs from parasitic nematodes. The phylogenies reveal 13 putative FLP-GPCRs that are highly conserved in nematode parasites (see Fig. 3). Entries are identified as per previous designation [28] and prefixes which indicate species: Ts, denotes *Trichinella spiralis*; Tm, denotes *Trichuris muris*; Bm, denotes *Brugia malayi*; Wb, denotes *Wuchereria bancrofti*; Di, denotes *Dirofilaria immitis*; Ov, denotes *Onchocerca volvulus*; Oo, denotes *Onchocerca ochengi*; Ll, denotes *Loa loa*; As, denotes *Ascaris suum*; Ac, denotes *Ancylostoma caninum*; Nb, denotes *Nippostrongylus brasiliensis*; Hc, denotes *Haemonchus contortus*; Ce, denotes *Caenorhabditis elegans*; Sr, denotes *Strongyloides ratti*; Bx, denotes *Bursaphelenchus xylophilus*; Gp, denotes *Globodera pallida*; Mh, denotes *Meloidogyne hapla*; Mi, denotes *Meloidogyne incognita*. Each phylogeny (A–C) represents a single *C. elegans* FLP-GPCR cluster identified in Fig. 3: (A) denotes the phylogeny of parasitic nematode FLP-GPCR homologues of the *C. elegans* predicted proteins present in Fig. 3 cluster 1; (B) denotes the phylogeny of parasitic nematode FLP-GPCR homologues of the *C. elegans* predicted proteins present in Fig. 3 cluster 2; (C) denotes the phylogeny of parasitic nematode FLP-GPCR homologues of the *C. elegans* predicted proteins present in Fig. 3 cluster 3. FLP-GPCR cluster nodes supported by a bootstrap analysis value of >70% are identified by red dots. Clusters are shaded and named according to the relevant *C. elegans* orthologue. Bootstrap values are shown as percentages. †Note that NPR-2, NPR-5, NPR-8, DMSR-1 and FRPR-18 isoforms were not delineated in the parasite species. In addition only the longest *C. elegans* FLP-/putative FLP-receptor isoforms were included in the phylogenetic analyses such that Ce_NPR-2 represents isoform a; Ce_NPR-5 represents isoform b; Ce_NPR-8 represents isoform b, Ce_DMSR-1 represents isoform a; Ce_EGL-6 represents isoform a; Ce_FRPR-18 represents isoform b.

Fig. 3; Supplementary Figure 2). Whilst the clustering does not necessarily indicate that closely related receptors will be activated by highly similar ligands, it is interesting to note that NPR-4 and NPR-10 cluster with a bootstrap value of 99% and are both activated by FLP-18 peptides. Also, cluster 1 (NPY-like receptors) displayed the highest complement of deorphanised FLP-GPCRs (eight), which provides an indication that those orphan receptors in this cluster may be activated by FLPs (see [50]). In contrast, NPR-22 (activated by *flp-7* encoded peptides) did not cluster strongly enough with any orphan *C. elegans* GPCRs to permit the designation of these related orphans as putative FLP-receptors.

3.2.3. Novel putative FLP-GPCR complement in parasitic nematodes

Predicted orphan *C. elegans* GPCR sequences that clustered with deorphanised *C. elegans* FLP-receptors (see Fig. 3) were used as search string queries in a BLAST approach to identify putative *flp*-GPCR orthologues in parasitic nematodes. Subsequently the predicted FLP-GPCRs (*C. elegans* and parasite species; containing >5 TM domains) were subjected to phylogenetic analysis (Fig. 4). Members of the Sex Peptide related receptor gene family (SPRR-1 and SPRR-2), present within cluster 2 (see Fig. 3), were excluded from this analysis as they are known homologues of a *Drosophila* receptor which is activated by a non-FLP neuropeptide [51]. Homologues of 13 putative *C. elegans flp*-GPCRs were identified in parasitic nematodes. The bootstrap values (>70%) provided by the phylogenetic analysis supported the designation of the majority (90.3%) of these putative parasite *flp*-GPCRs as orthologues, and validated our BLAST-based approach. Approximately 50% of the putative *flp*-GPCR homologues identified in the parasites were derived from *C. elegans* NPY-like receptors (see Fig. 3, cluster 1; Fig. 4A), eight of which have been deorphanised. Therefore, this may strengthen the prediction that these orphan parasite receptors are the most likely to have FLPs as ligands (see [50]) and may provide the primary focus for future deorphanisation efforts.

With respect to broad-spectrum drug target appeal, DMSR-2, -7 and -8 are the most highly conserved putative *C. elegans* FLP-GPCRs found in parasitic nematodes. It will be interesting to see if deorphanisation studies link these to highly conserved *flp*-ligands (e.g. *flp-1* and *flp-14* encoded peptides which remain unmatched with their cognate receptors). It is likely that those putative *C. elegans flp*-GPCRs not identified in the parasites (*npr-7*, *dmsr-9*, *dmsr-10*, *dmsr-11*, *dmsr-12*, *dmsr-13*, *dmsr-14*, *dmsr-15*, *dmsr-16*, *frpr-4*, *frpr-6*, *frpr-11*, *frpr-12*, and *frpr-16*) have evolved via gene duplication following the separation of the *Caenorhabditis* spp. and the parasites from their common ancestor.

Several putative FLP-GPCRs in the parasite species did not fall into orthologous clusters [see Fig. 4; e.g. sequences previously designated as NPR-10, DMSR-7 and FRPR-18 homologues in clade 2 species (*T. spiralis* and *T. muris*), and DMSR-4, and DMSR-5, in the plant parasitic nematodes (*B. xylophilus*, *M. incognita*, *M. hapla*, and *G. pallida*)]. In addition, several putative FLP-GPCRs designated as either NPR-1 or NPR-2 (*Ov*-NPR-1, *Oo*-NPR-1, *Di*-NPR-1, *Ll*-NPR-1, *As*-NPR-2, *Nb*-NPR-2.2, *Sr*-NPR-1.2, *Hc*-NPR-1 and *Nb*-NPR-1) failed to cluster with either *Ce*-NPR-1 or *Ce*-NPR-2 (see Fig. 5A) and as such constitute two NPR-1/2-like parasite specific clusters. All sequences designated

as either NPR-1 or NPR-2 homologues in this study exhibit a phenylalanine at position 215, linked to sociality in *C. elegans*.

4. Conclusions

This study provides the first pan-phylum genome-based overview of the FLPergic complement in parasitic nematodes. This encompasses both an update of the FLP ligand profile and reports, for the first time, the putative FLP-GPCR complement beyond *C. elegans*. Our study revises the number of *flp*-genes from 34 to 32, of which only one (*flp-31*) appears to be parasite-specific. This study also reveals that whilst the *flp*-signatures identified in nematode parasites are structurally similar to those described for *C. elegans*, the diversity in the *flp*-complement in the parasites is variably restricted. Indeed, nematode parasites only possess a proportion of the *C. elegans flp*-gene complement and exhibit inter-clade variation where clade-matched species broadly display similar *flp*-gene profiles.

With respect to FLP-GPCRs we have reported the conservation of known deorphanised *C. elegans* FLP-receptors in parasite species and our phylogenetic approach has facilitated the identification of a further 13 putative *flp*-GPCRs in nematode parasites. These data reveal that there may be many more nematode FLP-receptors than previously thought.

This dataset contributes significantly to future FLP-receptor deorphanisation efforts by providing a more comprehensive library of parasite-specific FLP ligands and a catalogue of putative parasite FLP-receptors against which to screen. These data help facilitate the functional characterisation of parasite FLP-GPCRs to reveal those most appealing for exploitation as novel chemotherapeutic targets.

Transparency document

The [Transparency document](#) associated with this article can be found in the online version.

Acknowledgements

This work was funded by the Biotechnology and Biological Sciences Research Council (BB/H019472/1) and the Department of Education and Learning for Northern Ireland. The funders had no role in study design, execution, or preparation of the manuscript.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at [doi:10.1016/j.euprot.2014.04.002](https://doi.org/10.1016/j.euprot.2014.04.002).

REFERENCES

- [1] Li J, Todd TC, Lee J, Trick HN. Biotechnological application of functional genomics towards plant-parasitic nematode control. *Plant Biotechnol J* 2011;9:936–44.

- [2] Waller PJ. From discovery to development: current industry perspectives for the development of novel methods of helminth control in livestock. *Vet Parasitol* 2006;139:1–14.
- [3] Prichard RK, Basáñez M-G, Boatin BA, McCarthy JS, Garcia HH, Yang G-J, et al. A research agenda for helminth diseases of humans: intervention for control and elimination. *PLoS Negl Trop Dis* 2012;6:e1549.
- [4] Epe C, Kaminsky R. New advancement in anthelmintic drugs in veterinary medicine. *Trends Parasitol* 2013;29:129–34.
- [5] Xiao S-H, Hui-Ming W, Tanner M, Utzinger J, Chong W. Tribendimidine: a promising, safe and broad-spectrum anthelmintic agent from China. *Acta Trop* 2005;94:1–14.
- [6] Hotez PJ. Mass drug administration and integrated control for the world's high-prevalence neglected tropical diseases. *Clin Pharmacol Ther* 2009;85:659–64.
- [7] Lustigman S, Prichard RK, Gazzinelli A, Grant WN, Boatin Ba, McCarthy JS, et al. A research agenda for helminth diseases of humans: the problem of helminthiasis. *PLoS Negl Trop Dis* 2012;6:e1582.
- [8] Jones JT, Haegeman A, Danchin EGJ, Gaur HS, Helder J, Jones MGK, et al. Top 10 plant-parasitic nematodes in molecular plant pathology. *Mol Plant Pathol* 2013;14:946–61.
- [9] McVeigh P, Atkinson L, Marks NJ, Mousley A, Dalzell JJ, Sluder A, et al. Parasite neuropeptide biology: seeding rational drug target selection? *Int J Parasitol Drugs Drug Resist* 2012;2:76–91.
- [10] McVeigh P, Geary TG, Marks NJ, Maule AG. The FLP-side of nematodes. *Trends Parasitol* 2006;22:385–96.
- [11] Li C, Kim K. Neuropeptides. *WormBook*; 2008. p. 1–36.
- [12] Ringstad N, Horvitz HR. FMRFamide neuropeptides and acetylcholine synergistically inhibit egg-laying by *C. elegans*. *Nat Neurosci* 2008;11:1168–76.
- [13] Larsen MJ, Lancheros ER, Williams T, Lowery DE, Geary TG, Kubiak TM. Functional expression and characterization of the *C. elegans* G-protein-coupled FLP-2 Receptor (T19F4.1) in mammalian cells and yeast. *Int J Parasitol Drugs Drug Resist* 2013;3:1–7.
- [14] Lowery DE, Geary TG, Kubiak TM, Larsen MJ. Pharmacia & Upjohn Company, G-protein coupled receptor-like receptors and modulators thereof, United States Patent 6,632,631 (2003).
- [15] Kubiak TM, Larsen MJ, Zantello MR, Bowman JW, Nulf SC, Lowery DE. Functional annotation of the putative orphan *Caenorhabditis elegans* G-protein-coupled receptor C10C6.2 as a FLP15 peptide receptor. *J Biol Chem* 2003;278:42115–20.
- [16] Kubiak TM, Larsen MJ, Nulf SC, Zantello MR, Burton KJ, Bowman JW, et al. Differential activation of social and solitary variants of the *Caenorhabditis elegans* G protein-coupled receptor NPR-1 by its cognate ligand AF9. *J Biol Chem* 2003;278:33724–9.
- [17] Kubiak TM, Larsen MJ, Bowman JW, Geary TG, Lowery DE. FMRFamide-like peptides encoded on the flp-18 precursor gene activate two isoforms of the orphan *Caenorhabditis elegans* G-protein-coupled receptor Y58G8A.4 heterologously expressed in mammalian cells. *Biopolymers* 2008;90:339–48.
- [18] Keating CD, Kriek N, Daniels M, Ashcroft NR, Hopper NA, Siney EJ, et al. Whole-genome analysis of 60 G protein-coupled receptors in *Caenorhabditis elegans* by gene knockout with RNAi. *Curr Biol* 2003;13:1715–20.
- [19] Mertens I, Meeusen T, Janssen T, Nachman R, Schoofs L. Molecular characterization of two G protein-coupled receptor splice variants as FLP2 receptors in *Caenorhabditis elegans*. *Biochem Biophys Res Commun* 2005;330:967–74.
- [20] Mertens I, Clinckspoor I, Janssen T, Nachman R, Schoofs L. FMRFamide related peptide ligands activate the *Caenorhabditis elegans* orphan GPCR Y59H11AL.1. *Peptides* 2006;27:1291–6.
- [21] McVeigh P, Leech S, Mair GR, Marks NJ, Geary TG, Maule AG. Analysis of FMRFamide-like peptide (FLP) diversity in phylum Nematoda. *Int J Parasitol* 2005;35:1043–60.
- [22] Elsworth B, Wasmuth J, Blaxter M. NEMBASE4: the nematode transcriptome resource. *Int J Parasitol* 2011;41:881–94.
- [23] Holterman M, van der Wurff A, van den Elsen S, van Megen H, Bongers T, Holovachov O, et al. Phylum-wide analysis of SSU rDNA reveals deep phylogenetic relationships among nematodes and accelerated evolution toward crown Clades. *Mol Biol Evol* 2006;23:1792–800.
- [24] Laing R, Kikuchi T, Martinelli A, Tsai IJ, Beech RN, Redman E, et al. The genome and transcriptome of *Haemonchus contortus*, a key model parasite for drug and vaccine discovery. *Genome Biol* 2013;14:R88.
- [25] Schwarz EM, Korhonen PK, Campbell BE, Young ND, Jex AR, Jabbar A, et al. The genome and developmental transcriptome of the strongylid nematode *Haemonchus contortus*. *Genome Biol* 2013;14:R89.
- [26] Mertens I, Vandingenen A, Meeusen T, Janssen T, Luyten W, Nachman RJ, et al. Functional characterization of the putative orphan neuropeptide G-protein coupled receptor C26F1.6 in *Caenorhabditis elegans*. *FEBS Lett* 2004;573:55–60.
- [27] Rogers C, Reale V, Kim K, Chatwin H, Li C, Evans P, et al. Inhibition of *Caenorhabditis elegans* social feeding by FMRFamide-related peptide activation of NPR-1. *Nat Neurosci* 2003;6:1178–85.
- [28] Froominckx L, Van Rompay L, Temmerman L, Van Sinay E, Beets I, Janssen T, et al. Neuropeptide GPCRs in *C. elegans*. *Front Endocrinol (Lausanne)* 2012;3:167.
- [29] Kikuchi T, Cotton Ja, Dalzell JJ, Hasegawa K, Kanzaki N, McVeigh P, et al. Genomic insights into the origin of parasitism in the emerging plant pathogen *Bursaphelenchus xylophilus*. *PLoS Pathog* 2011;7:e1002219.
- [30] Abad P, Gouzy J, Aury J-M, Castagnone-Sereno P, Danchin EGJ, Deleury E, et al. Genome sequence of the metazoan plant-parasitic nematode *Meloidogyne incognita*. *Nat Biotechnol* 2008;26:909–15.
- [31] Dalzell JJ, McVeigh P, Warnock ND, Mitreva M, Bird DM, Abad P, et al. RNAi effector diversity in nematodes. *PLoS Negl Trop Dis* 2011;5:e1176.
- [32] Lu G. Software review Vector NTI, a balanced all-in-one sequence analysis suite. *Brief Bioinform* 2004;5:378–88.
- [33] Tusnády GE, Simon I. The HMMTOP transmembrane topology prediction server. *Bioinformatics* 2001;17:849–50.
- [34] Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol* 2011;28:2731–9.
- [35] Thompson JD, Higgins DG, Gibson TJ. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res* 1994;22:4673–80.
- [36] Zamanian M, Kimber MJ, McVeigh P, Carlson Sa, Day TA. The repertoire of G protein-coupled receptors in the human parasite *Schistosoma mansoni* and the model organism *Schmidtea mediterranea*. *BMC Genomics* 2011;12:596.
- [37] Jones DT, Taylor WR, Thornton JM. The rapid generation of mutation data matrices from protein sequences. *Comput Appl Biosci* 1992;8:275–82.
- [38] Saitou N, Nei M. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* 1987;4:406–25.
- [39] Jarecki JL, Frey BL, Smith LM, Stretton AO. Discovery of neuropeptides in the nematode *Ascaris suum* by database mining and tandem mass spectrometry. *J Proteome Res* 2011;10:3098–106.

- [40] Atkinson LE, Stevenson M, McCoy CJ, Marks NJ, Fleming C, Zamanian M, et al. flp-32 Ligand/receptor silencing phenocopy faster plant pathogenic nematodes. *PLoS Pathog* 2013;9:e1003169.
- [41] Maule AG, Mousley A, Marks NJ, Day TA, Thompson DP, Geary TG, et al. Neuropeptide signaling systems – potential drug targets for parasite and pest control. *Curr Top Med Chem* 2002;2:733–58.
- [42] Mitreva M, Jasmer DP, Zarlenga DS, Wang Z, Abubucker S, Martin J, et al. The draft genome of the parasitic nematode *Trichinella spiralis*. *Nat Genet* 2011;43:228–35.
- [43] Jarecki JL, Andersen KM, Miller AH, Ramaker MA, Vestling MM, Stretton AO. Three independent techniques localize expression of transcript of p-11 and its bioactive peptide products to the paired AVK neurons in *Ascaris suum*: in situ hybridization, immunocytochemistry, and single cell mass spectrometry. *ASC Chem Neurosci* 2013;4:418–34.
- [44] McVeigh P, Mair GR, Atkinson L, Ladurner P, Zamanian M, Novozhilova E, et al. Discovery of multiple neuropeptide families in the phylum Platyhelminthes. *Int J Parasitol* 2009;39:1243–52.
- [45] Cohen M, Reale V, Olofsson B, Knights A, Evans P, de Bono M. Coordinated regulation of foraging and metabolism in *C. elegans* by RFamide neuropeptide signaling. *Cell Metab* 2009;9:375–85.
- [46] Janssen T, Lindemans M, Meelkop E, Temmerman L, Schoofs L. Coevolution of neuropeptidergic signaling systems: from worm to man. *Ann N Y Acad Sci* 2010;1200:1–14.
- [47] Chalasani SH, Kato S, Albrecht DR, Nakagawa T, Abbott LF, Bargmann CI. Neuropeptide feedback modifies odor-evoked dynamics in *Caenorhabditis elegans* olfactory neurons. *Nat Neurosci* 2010;13:615–21.
- [48] De Bono M, Bargmann CI. Natural variation in a neuropeptide Y receptor homolog modifies social behavior and food response in *C. elegans*. *Cell* 1998;94:679–89.
- [49] Caers J, Verlinden H, Zels S, Vandersmissen HP, Vuerinckx K, Schoofs L. More than two decades of research on insect neuropeptide GPCRs: an overview. *Front Endocrinol (Lausanne)* 2012;3:151.
- [50] Vogel KJ, Brown MR, Strand MR. Phylogenetic investigation of Peptide hormone and growth factor receptors in five dipteran genomes. *Front Endocrinol (Lausanne)* 2013;4: 193.
- [51] Yapici N, Kim Y-J, Ribeiro C, Dickson BJ. A receptor that mediates the post-mating switch in *Drosophila* reproductive behaviour. *Nature* 2008;451:33–7.